

BEST AVAILABLE COPY

PCT / IB 04 / 03806
22 NOV 2004

PA 1223634

REC'D 22 NOV 2004

WIPO

PCT

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

September 14, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/526,681

FILING DATE: December 03, 2003

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



P. SWAIN
Certifying Officer

Please type a plus sign (+) inside this box ☐

Approved for use through 04/30/2003. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PTO/SB/16 (5-03)
22386 U.S. PTO
60/526681



PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
Tonia Sue Everett Lee Frederick Henry		Agin Rosey Weber		Schoolcraft, Michigan Schoolcraft, Michigan Mattawan, Michigan	
<input type="checkbox"/> Additional inventors are being named on the _____ separately numbered sheets attached hereto					
TITLE OF THE INVENTION (280 characters max)					
IN OVO VACCINATION OF CAMPYLOBACTER IN AVIAN SPECIES					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input checked="" type="checkbox"/> Customer Number		23389		<div>Place Customer Number Bar Code Label here</div>	
OR Type Customer Number here					
<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		ZIP	
Country		Telephone		Fax	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification		Number of Pages		11	
<input type="checkbox"/> Drawing(s)		Number of Sheets		None	
<input type="checkbox"/> Application Data Sheet		See 37 CFR 1.76			
<input type="checkbox"/> CD(s)		Number			
<input type="checkbox"/> Other (specify)					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)					
<input checked="" type="checkbox"/> A check or money order is enclosed to cover the filing fees				FILING FEE AMOUNT (\$)	
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number		19-1013/SSMP		\$160.00	
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.					
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

Respectfully submitted,

SIGNATURE

TYPED or PRINTED NAME Peter I. Bernstein

(516) 742-4343

TELEPHONE

Date

12/3/03

REGISTRATION NO.

43,497

(if appropriate)

Docket Number:

P16171(PC25353)

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is used by the public to file (and by the PTO to process) a provisional application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the complete provisional application to the PTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

P19LARGE/REV05

CERTIFICATE OF MAILING BY "EXPRESS MAIL" (37 CFR 1.10)

Applicant(s): Tonia Sue Agin, et al.

Docket No.

P16171(PC25353)

Serial No.
To Be AssignedFiling Date
HerewithExaminer
UnassignedGroup Art Unit
UnassignedInvention: **IN OVO VACCINATION OF CAMPYLOBACTER IN AVIAN SPECIES**I hereby certify that this New Provisional Patent Application

(Identify type of correspondence)

is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 in an envelope addressed to: Director of the United States Patent and Trademark Office, P.O. Box

1450, Alexandria, VA 22313-1450 on

December 3, 2003

(Date)

Peter I. Bernstein

(Typed or Printed Name)


(Signature)EV247990382US

("Express Mail" Mailing Label Number)

Note: Each paper must have its own certificate of mailing.

In Ovo Vaccination Of Campylobacter In Avian Species

5

FIELD OF THE INVENTION

This invention relates to methods of inducing an immune response in an avian species against *Campylobacter* by administering, *in ovo*, live *Campylobacter* cells.

10

BACKGROUND OF THE INVENTION

Consumption of poultry contaminated with *Campylobacters* has been implicated as a major source of human infection. Therefore, removal of these organisms from the poultry food chain has been a significant objective of *Campylobacter* research.

15

Campylobacters typically grow and colonize in the avian gut environment. Colonization has been reported in chickens, ducks, pigeons, quail, ostriches and turkeys. Colonization of poultry does not result in disease but is commensal in nature.

20

The use of a competitive exclusion culture to exclude *Salmonella* or *Campylobacter* from the digestive tract of a bird has been described in U.S. Patent No. 6,491,910. The efficacy of this method against *Campylobacter* appears variable. *In ovo* vaccination with heat killed cells of *Campylobacter jejuni* has been described by Noor et al. (*British Poultry Science*, 1995, 36(4): 563-73) and by Noor (*Jurnal Ilmu Ternak Dan Veteriner*, 1998, 3(4): 264-269). *In ovo* immunization of chickens with flagellin and whole cell protein antigens of *Campylobacter jejuni* has also been reported by S. Noor et al. (*Jurnal Ilmu Ternak Dan Veteriner*, 2000, 5(2): 119-124). Efforts have also been made in order to identify the genes involved in colonization (Ziprin et al., Abstracts, Poultry Science Association meeting, August 8-11, 1999, Springdale, AR). See also, Rice, 1997, "Campylobacter jejuni in broiler chickens: colonization and humoral immunity following oral vaccination and experimental infection", *Vaccine* 15(17-18): 1922-1932, wherein killed *Campylobacter* cells are administered; and Ziprin et al, 2002, *Current Microbiology* 44(3): 221-223, wherein chicks were vaccinated post-hatch with viable but non-colonizing strains.

25

30

Prior to the present invention, there has been no effective immunization strategy that employs *in ovo* administration with live *Campylobacter* cells.

35

SUMMARY OF THE INVENTION

The present invention provides a method of inducing an immune response in an avian species against *Campylobacter* by administering, *in ovo*, live cells of *Campylobacter*.

According to the present invention, live cells of *Campylobacter* can be safely administered to any avian species for the purpose of inducing an immune response against *Campylobacter*, especially a domesticated avian species such as chicken, turkey, duck, goose and quail.

Campylobacter species suitable for use in the present method include, but are not limited to, *Campylobacter jejuni*, *Campylobacter coli*, *Campylobacter lari*, or a combination thereof.

The cells of *Campylobacter* can be wild type cells or cells of *Campylobacter* that have been genetically modified to contain one or more mutations in the genome, or to contain a desirable heterologous sequence.

Preferably, the cells are combined with a veterinary-acceptable carrier prior to administration and are administered in an amount that is effective to induce an immune response in the avian species developed from the treated egg, preferably in an amount of at least about 5×10^5 , and more preferably, of at least about 1×10^6 live cells.

DETAILED DESCRIPTION OF THE INVENTION

It has been surprisingly found in accordance with the present invention that live cells of *Campylobacter* can be safely administered *in ovo*, to eggs of an avian species, which results in effective colonization and induces an immune response in the avian species against *Campylobacter*.

Accordingly, the present invention provides a method of inducing an immune response in an avian species against *Campylobacter* by *in ovo* administration of live cells of *Campylobacter*.

As used herein, the term "avian species" or "bird" is meant to include any avian species, including a domestic or a game bird, e.g., chicken, turkey, duck, goose, or quail. Preferably, live cells of *Campylobacter* are administered to eggs of a domesticated bird raised for commercial production of eggs or meat, such as a domesticated chicken, turkey, duck, goose and quail.

The term "*Campylobacter*" refers to any *Campylobacter* species or any strain of a *Campylobacter* species, including *Campylobacter jejuni*, *Campylobacter coli*, *Campylobacter lari*. *Campylobacter* strains suitable for use in the present method include *C. jejuni* UA535 and *C. jejuni* 81-176, as well as the following mutant strains: *C. jejuni* CsrA (mutated within gene Cj1103, a *csrA* homolog), *C. jejuni* HspR (Cj1230), *C. jejuni* HtrA (Cj1228c), *C. jejuni* Dps (Cj1534c), *C. jejuni* flbA (Cj0822c), *C. jejuni* Pnp (Cj1253), and *C. jejuni* SurE (Cj0293).

Both wild type and mutant strains of a *Campylobacter* species can be used in the method of the present invention, including strains that have been genetically engineered and contain one or more mutations in the genome. The capacity of a mutant *Campylobacter* strain in colonization and induction of an immune response in an avian species can be determined following the techniques described in the examples hereinbelow or techniques known to those skilled in the art.

In a preferred embodiment, the *Campylobacter* strain employed in the method of the present invention has been genetically engineered to contain a heterologous polynucleotide sequence. The heterologous sequence can be transferred into the *Campylobacter* strain via a plasmid, phage or cosmid vector by various means such as conjugation, electroporation, or transformation. Other methods such as transduction are also suitable, wherein the recombinant DNA in the form of a transducing phage or cosmid vector is packaged within a phage. Once the recombinant polynucleotide is in the carrier *Campylobacter* strain, it may continue to exist as a separate autonomous replicon or it may insert into the *Campylobacter* chromosome and be reproduced along with the chromosome during cell division.

According to the present invention, the heterologous polynucleotide sequence can encode an antigen from an organism, such as a virus, bacterium or parasite, that causes disease in bird or that causes food-borne illness in humans upon consumption of a bird contaminated with the organism. Administration of live cells of genetically engineered *Campylobacter* containing such heterologous sequence can induce an immune response in the avian species against both *Campylobacter* and the pathogenic organism. Examples of such pathogenic organisms include *Salmonella*, *Escherichia coli*, *Eimeria*, *Clostridium*, infectious bursal disease virus.

The heterologous polynucleotide sequence can also encode a protein essential in colonization of domesticated birds by *Campylobacter*. Proteins known to be essential in colonization of *Campylobacter* include, e.g., the gene products of *dnaJ* and *cadF*.

Additionally, the heterologous polynucleotide sequence can encode a protein or peptide that stimulates the immune system of bird. Examples of proteins or peptides that stimulate the immune system of birds include, e.g., cholera toxin or *E. coli* heat labile toxin.

Moreover, the heterologous polynucleotide sequence can itself enhance the growth or feed efficiency of a domesticated bird, or can encode a protein or peptide that enhances the growth or feed efficiency of a domesticated bird. Examples of such molecules or proteins include epidermal growth factor, insulin-like growth factor, interleukins, and antimicrobial peptides.

According to the present invention, a *Campylobacter* species is cultured by standard methods known to those skilled in the art and live cells of such *Campylobacter* are collected for *in ovo* administration to eggs of an avian species. Live cells of more than one *Campylobacter* species can be combined for *in ovo* administration.

In addition to *Campylobacter* cells, a veterinary-acceptable carrier can also administered. Preferably, *Campylobacter* cells and a veterinary-acceptable carrier are both administered *in ovo*, either together or separately. Alternatively, a veterinary-acceptable carrier can be administered to the bird any time post-hatch in feed or water, or by aerosol spray. The live *Campylobacter* cells administered, whether with or without a veterinary-acceptable carrier, are free of neutralizing

factors, such as neutralizing antibodies or fragments of antibodies, as described in U.S. Patent 6,440,408.

The term "a veterinary-acceptable carrier" includes solvents, dispersion media, coatings, adjuvants, stabilizing agents, diluents, preservatives, antifungal agents, isotonic agents, adsorption delaying agents, and the like. Diluents can include water, saline, dextrose, ethanol, glycerol, and the like. Isotonic agents can include sodium chloride, dextrose, mannitol, sorbitol, and lactose, among others. Stabilizers include albumin, among others.

Adjuvants suitable for use in the present method include but are not limited to: mineral gels, e.g., aluminum hydroxide; surface active substances such as lysolecithin; glycosides, e.g., saponin derivatives such as Quil A or GPI-0100 (United States Patent No. 5,977,081); cationic surfactants such as DDA, pluronic polyols; polyanions; non-ionic block polymers, e.g., Pluronic F-127 (B.A.S.F., USA); peptides; mineral oils, e.g. Montanide ISA-50 (Seppic, Paris, France), carbopol, Amphigen (Hydronics, Omaha, NE USA), Alhydrogel (Superfos Biosector, Frederikssund, Denmark) oil emulsions, e.g. an emulsion of mineral oil such as BayolF/Arlacel A and water, or an emulsion of vegetable oil, water and an emulsifier such as lecithin; alum, cholesterol, rmlT, cytokines and combinations thereof. The immunogenic component may also be incorporated into liposomes, or conjugated to polysaccharides and/or other polymers for use in a vaccine formulation. Additional substances that can be included in a product for use in the present methods include, but are not limited to one or more preservatives such as disodium or tetrasodium salt of ethylenediaminetetracetic acid (EDTA), merthiolate, and the like. Immunostimulants which enhance the immune system's response to antigens may also be included in a product. Examples of suitable immunostimulants include cytokines such as IL-12 or IL-2, or stimulatory molecules such as muramyl dipeptide, aminoquinolones, lipopolysaccharide, and the like. The adjuvant can be combined with live cells of *Campylobacter* prior to *in ovo* administration, or can be administered independently any time after hatching by way of feed, water or aerosol spray, provided to the bird.

The live cells are administered at a dose effective to induce an immune response against *Campylobacter* in the avian species developed from the treated egg. The amount of live cells of *Campylobacter* that is effective to induce an immune response, or "the immunizing effective amount", may vary, depending on the particular species or strains of *Campylobacter* used in the administration and the species of bird being immunized. Generally speaking, at least about 5×10^5 live cells should be administered to be effective in inducing an immune response. More preferably, at least about 1×10^6 live cells of *Campylobacter* are administered.

By "inducing an immune response" is meant that the live cells of *Campylobacter* administered to an egg induce an immune response in the bird developed from the egg, which in turn provides some degree of protection to the bird against colonization of a *Campylobacter*

species, which can be the same as, or different from, the *Campylobacter* species used in the *in ovo* administration.

An immune response induced by *in ovo* administration of live cells of *Campylobacter* can be a cellular immune response mediated primarily by cytotoxic T-cells, or a humoral immune response mediated primarily by helper T-cells, which in turn activates B-cells leading to antibody production, or the combination of a cellular immune response and a humoral immune response.

According to the present invention, one or more additional immunogens can be included in the administration *in ovo*. Such immunogens include antigens derived from viruses, e.g. avian infectious bronchitis virus, avian infectious bursal disease virus, avian encephalomyelitis virus, egg drop syndrome virus, influenza virus, reovirus, adenovirus, hydropericardium syndrome virus, among others; antigens derived from bacteria, e.g., *Haemophilus paragallinarum*, *Salmonella typhimurium*, *S. enteritidis*, *S. pullori*, *S. gallinarum*, *S. choleraesuis*, *E. coli*, *Clostridium spp.*, *Mycoplasma spp.*, *enterococcus*, among others; and antigens derived from protozoan, e.g. *Eimeria tenella*, *E. maxima*, *E. acervulina*, *E. brunetti*, *E. necatrix*, among others.

By "*in ovo* administration" is meant administration to eggs of an avian species, preferably eggs in the fourth quarter of incubation. That is, for chicken eggs, the administration is conducted preferably on about the fifteenth to nineteenth day of incubation, and more preferably on about the eighteenth day of incubation. For turkey eggs, the administration is conducted preferably on about the twenty-first to twenty-sixth day of incubation, and more preferably on about the twenty-fifth day of incubation.

The administration can be conducted by any method which results in the introduction of live cells of *Campylobacter* into an egg through the shell. A preferred method of administration is by injection. The injection can be made at any site of an egg, so long as the injection does not damage the tissues or organs of the embryo, or the extraembryonic membranes surrounding the embryo. The injection can be achieved by using any one of the well-known egg injection devices, such as a conventional hypodermic syringe fitted with a needle of about 18 to 22 gauge, or a high speed automated egg injection system as described in U.S. Patent No. 4,681,063, 4,040,388, 4,469,047, and 4,593,646.

The present invention is further illustrated by the following non-limiting examples.

EXAMPLE 1

Broiler chicken eggs were injected *in ovo* on day 18 of incubation with 10^6 colony forming units (CFU) of *Campylobacter jejuni* UA535. This is a wild type, unmodified strain of *C. jejuni*, originally isolated from a human clinical case of campylobacteriosis. A second group of eggs remained uninoculated until the day of hatch, at which time the chicks in this group were inoculated *per os* with 10^6 CFU *C. jejuni* UA535. At hatch, the number of live chicks and

unhatched eggs in each treatment were recorded and all chicks were placed on litter with 3 replicate pens of twenty birds per treatment.

In ovo inoculation had no impact on hatch rate, since 97.5% of uninoculated eggs and 98.8% of inoculated eggs hatched. Both groups were similar in weight at hatch and gained weight at a similar rate throughout the study. The two routes of administration of *C. jejuni* to broiler chicks resulted in equivalent colonization both in terms of the number of birds colonized and the number of *C. jejuni* per gram of cecal contents. No signs of clinical significance were observed in any of the birds. These data indicate that *in ovo* inoculation of broiler chicks with *C. jejuni* is as safe as *per os* inoculation.

The geometric mean antibody titer in response to *in ovo* inoculation was at least as great as, if not greater than, the titer resulting from *per os* inoculation at day of hatch. Therefore, not only can wild type *C. jejuni* UA535 be safely administered *in ovo* to broiler chicks, but administration by this route results in robust colonization and immune response.

EXAMPLE 2

To demonstrate that colonization after *in ovo* inoculation with *C. jejuni* is not unique to strain UA535, colonization of birds by another human clinical isolate, *C. jejuni* 81-176, was evaluated. Groups of broiler eggs were injected on Day 18 of embryo incubation with 1.8×10^5 or 1.4×10^7 CFU of *C. jejuni* 81-176. A third group of eggs remained uninoculated. At hatch, a group of chicks from the uninoculated eggs were retained as uninoculated control birds, and a second group of chicks from these eggs were given 8.3×10^6 CFU of *C. jejuni* 81-176 orally.

The data set obtained demonstrated that *in ovo* administration of *C. jejuni* strain 81-176 is at least as effective at colonizing broiler chicks as oral administration on the day of hatch.

EXAMPLE 3

On day 18 of embryo incubation, a group of broiler chicken eggs were inoculated *in ovo* with 7×10^7 CFU of *C. jejuni* CsrA. This strain has a mutation in CsrA, a protein involved in global regulation of carbon storage. A second group of eggs were retained as uninoculated controls. At hatch, a group of chicks from the uninoculated eggs were given 3×10^8 CFU of *C. jejuni* CsrA orally. Another group of chicks hatched from uninoculated eggs were retained as controls.

The results indicated that the mutant *C. jejuni* strain colonized all of the birds when administered *in ovo*, yet colonized only 3 out of 6 birds when given orally. Also, the mutant was able to fully colonize birds and induce an antibody response when administered *in ovo*, but appeared to be less effective at colonizing birds when given by the oral route on the day of hatch.

EXAMPLE 4

On day 18 of embryo incubation, groups of broiler chicken eggs were inoculated *in ovo* with 9×10^6 CFU of *C. jejuni* HspR, 8×10^7 CFU of *C. jejuni* HtrA, or 6×10^7 CFU of *C. jejuni* Dps.

5 The HspR strain has a mutation in a heat shock protein; the HtrA strain has a mutation in a serine protease gene; and the Dps strain is mutated in a gene involved in iron acquisition. Another group of eggs was retained as uninoculated controls. At hatch, groups of chicks from the uninoculated eggs were given 4×10^6 CFU of *C. jejuni* HspR, 1×10^8 CFU of *C. jejuni* HtrA, or 1×10^8 CFU of *C. jejuni* Dps orally. Another group of chicks hatched from uninoculated eggs was retained as
10 controls.

The HspR mutant was undetectable in samples from birds inoculated *per os*, but half the birds given this strain *in ovo* were colonized with *C. jejuni*. The results indicate that this mutant was able to colonize broiler chicks after *in ovo* administration, but not after oral administration on the day of hatch. The other mutants readily colonized birds and induced an immunological
15 response regardless of route of administration.

EXAMPLE 5

On day 18 of embryo incubation, groups of broiler chicken eggs were inoculated *in ovo* with 7.3×10^7 CFU of *C. jejuni* flbA, 1.1×10^8 CFU of *C. jejuni* Pnp, or 1.2×10^8 CFU of *C. jejuni* SurE. The FlbA strain is mutated in a gene encoding for a flagellar structural protein; the Pnp strain is mutated in a gene encoding for a nucleotidyltransferase; and the SurE strain is mutated in a gene encoding a phosphatase. Another group of eggs was retained as uninoculated controls. At hatch, groups of chicks from the uninoculated eggs were given 9.3×10^7 CFU of *C. jejuni* flbA,
20 1.6×10^8 CFU of *C. jejuni* Pnp, or 8×10^7 CFU of *C. jejuni* SurE orally. Another group of chicks hatched from uninoculated eggs was retained as controls.

The FlbA mutant demonstrated a transient colonization when given by the *in ovo* route, and this mutant appeared unable to colonize by the oral route. The Pnp mutant colonized birds effectively regardless of route of administration. The SurE mutant colonized birds effectively when
25 administered by the *in ovo* route, but was unable to colonize birds when administered orally.
30

EXAMPLE 6

35 On Day 18 of incubation, a group of broiler chicken eggs were injected *in ovo* with 2×10^6 cells of wild-type *C. jejuni* strain UA535. A second group of eggs were maintained as uninoculated controls. At hatch, a group of birds from uninoculated eggs were inoculated *per os* with 2×10^6 cells of *C. jejuni* UA535. The remaining birds that hatched from uninoculated eggs were kept as

controls. At 27 days post-hatch, half of the birds in each group were placed on medicated water containing 100 mg/L kanamycin to eliminate the immunizing *C. jejuni* strain. These infected/cleared birds would therefore be considered "immunized".

At 34 days post-hatch, selected groups of birds were exposed to birds previously infected with *C. jejuni* CjM20, a strain genetically-modified to be resistant to kanamycin. This method allowed the spread of CjM20 from the infected birds to the "immunized" birds. By diagnostic means, it was possible to distinguish the strain CjM20 from the immunizing strain UA535 in the samples collected.

Non-immunized, non-medicated birds challenged with CjM20 were fully colonized by 41 days of age. Colonization was delayed somewhat by kanamycin treatment, with about half the birds colonized by Day 41. All of the non-immunized, medicated, challenged birds were fully colonized by 49 days of age. Both groups of immunized birds had reduced numbers of *C. jejuni* compared with their non-immunized counterparts. All recovered bacteria appeared resistant to kanamycin, indicating that the immunizing strain was eliminated by kanamycin treatment. Immunized, non-challenged birds remained free of *C. jejuni* throughout the study. Thus, immunization of broiler chicks against colonization with *C. jejuni* was at least as effective by the *in ovo* route as by the oral route.

WHAT IS CLAIMED IS:

1. A method of inducing an immune response in a bird against *Campylobacter*, comprising administering, *in ovo*, during the final quarter of incubation, an immunizing effective amount of live cells of a *Campylobacter* species.
2. The method of claim 1, wherein said bird is a domesticated bird.
3. The method of claim 2, wherein said domesticated bird is selected from the group consisting of a chicken, a turkey, and a duck.
4. The method of claim 1, wherein said species of *Campylobacter* used in the administration is selected from the group consisting of *C. jejuni*, *C. coli*, and *C. lari*.
5. The method of claim 1, wherein the live cells used in the administration comprise live cells of more than one species of *Campylobacter*.
6. The method of claim 1, wherein the live cells are wild type or have been modified genetically.
7. The method of claim 6, wherein a heterologous polynucleotide sequence has been introduced into the live cells of *Campylobacter*.
8. The method of claim 7, wherein said heterologous polynucleotide sequence encodes a protein essential in colonization of domesticated birds by *Campylobacter*.
9. The method of claim 7, wherein said heterologous polynucleotide sequence encodes an antigen from a virus, bacteria, or parasite that causes disease in a domesticated bird.
10. The method of claim 7, wherein said heterologous polynucleotide sequence encodes an antigen from an organism that causes food-borne illness in humans.
11. The method of claim 7, wherein said heterologous polynucleotide sequence encodes a protein that enhances the growth or feed efficiency of a domesticated bird.

12. The method of claim 7, wherein said heterologous polynucleotide sequence encodes a protein that stimulates the birds' immune system.

13. The method of claim 1, further comprising administering a veterinary-acceptable carrier.

5

14. The method of claim 13, wherein said veterinary-acceptable carrier is combined with the live cells of *Campylobacter* prior to *in ovo* administration.

15. The method of claim 13, wherein said veterinary-acceptable carrier is administered to the bird in feed or water, or by aerosol spray, at any time after hatching.

10

16. The method of claim 14 or 15, wherein said veterinary-acceptable carrier is an adjuvant.

17. The method of claim 14 or 15, wherein said adjuvant has an immune-stimulating activity.

15

18. The method of claim 1, wherein live cells of *Campylobacter* are combined with at least one other immunogen selected from a viral, a bacterial or a protozoan immunogen.

20

ABSTRACT

5

The present invention provides a method of inducing an immune response against *Campylobacter* in an avian species, especially a domesticated avian species such as chicken, turkey, duck, goose and quail, by administering, *in ovo*, live cells of *Campylobacter*.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.